

The effects of fermentation and enzymatic treatment of pea on nutrient digestibility and growth performance of broilers

F. Goodarzi Borojeni^{1†}, M. Senz², K. Kozłowski³, D. Boros⁴, M. Wisniewska⁴, D. Rose⁵, K. Männer¹ and J. Zentek¹

¹Department of Veterinary Medicine, Institute of Animal Nutrition, Freie Universität Berlin, Königin-Luise-Str. 49, 14195 Berlin, Germany; ²Department Bioprocess Engineering and Applied Microbiology, Research and Teaching Institute for Brewing in Berlin, Institute of Biotechnology and Water, Seestrasse 13, 13353 Berlin, Germany; ³Department of Poultry Science, Faculty of Animal Bioengineering, University of Warmia and Mazury in Olsztyn, Oczapowskiego 5, 10-719 Olsztyn, Poland; ⁴Laboratory of Quality Evaluation of Plant Materials, Institute of Plant Breeding and Acclimatization – National Research Institute, 05-870 Radzikow, Blonie, Poland; ⁵Department of Food Biotechnology and Food Process Engineering, Berlin University of Technology, Seestrasse 13, 13353 Berlin, Germany

(Received 11 October 2016; Accepted 30 January 2017; First published online 18 April 2017)

*The present study examined the impacts of native, fermented or enzymatically treated peas (*Pisum sativum* L.) inclusion in broiler diets, on growth performance and nutrient digestibility. For the fermentation process, Madonna pea was mixed with water (1/1) containing 2.57×10^8 *Bacillus subtilis* (GalliPro[®]) spores/kg pea and then, incubated for 48 h at 30 °C. For the enzymatic treatment process, the used water for dough production contained three enzymes, AlphaGal[™] (α -galactosidase), RONOZYME[®] ProAct and VP (protease and pectinases respectively – DSM, Switzerland) and the pea dough incubated for 24 h at 30 °C. Nine corn-wheat-soybean diets were formulated by supplying 10%, 20% and 30% of the required CP with either native, fermented or enzymatically treated peas. Performance was recorded weekly and at the end of the experiment (day 35), apparent ileal digestibility (AID) of CP, amino acids (AA), crude fat, starch, Ca, P and K were determined. Data were subjected to ANOVA using GLM procedure with a 3 × 3 factorial arrangement of treatments. Both processes reduced α -galactosides, phytate, trypsin inhibitor activity and resistant starch in peas. Increasing levels of pea products up to 300 g/kg diet, reduced BW gain and feed intake ($P \leq 0.05$). Broilers fed diets containing enzymatically treated pea had the best feed conversion ratio at day 35. Different types of pea product and their inclusion levels had no effect on AID of all nutrients. The interaction between type of the pea products and inclusion levels was significant for AID of starch. For native pea diets, 10% group showed similar AID of starch to 20% native pea but it had higher AID than 30% native pea. For fermented and enzymatically treated groups, all three levels displayed similar AID of starch. In conclusion, enzymatic treatment and fermentation could improve the nutritional quality of pea. Inclusion of enzymatically treated pea in broiler diets could improve broiler performance compared with other pea products while, it displayed neither positive nor negative impact on nutrient digestibility. The present findings indicate the feasibility of these processes, particularly enzymatic treatment, for improving the nutritional quality of pea as a protein source for broiler nutrition.*

Keywords: anti-nutritional factors, enzymatic treatment, legumes, solid state fermentation, α -galactosides

Implications

There is an urgent need for new plant proteins for poultry nutrition. As is, the majority of protein needs for poultry production are covered by the soybean products. The dependence of European feed production industry on soybean as well as the growing economic, environmental and ethical concerns on the loss of ecologically important rainforests have led to an increasing demand for home-grown and sustainable protein sources (e.g. European domestic grain legumes) for feeding livestock. The present

study aimed to evaluate innovative and classic biotechnological approaches for production of high quality alternatives for soybean meal from domestic European peas.

Introduction

Soybean meal (SBM) is the most common plant protein source in poultry nutrition. Environmental factors remarkably limit cultivation of soybean in many parts of the world, especially Europe. In Europe, domestic legumes like peas, beans and lupines or oilseeds like rapeseed can be alternative plant protein sources. Peas have relatively high CP. They also can potentially provide considerable

[†] E-mail: farshad.goodarzi@fu-berlin.de

amount of energy due to their high starch contents. However, peas contain also variable amounts of anti-nutritional factors (ANF) such as α -galactosides, trypsin inhibitors (TI), resistant starch (RS), pectin, tannins, lectin, phytic acid and non-starch polysaccharides (NSP), which can significantly impair the digestive process especially in young chicks. This may lead to pancreatic hypertrophy, increased nutritional requirement for sulfur-containing amino acids (AA) and finally poor bird performance (Igbasan and Guenter, 1996; Nalle *et al.*, 2011; Frikha *et al.*, 2013).

Different processes like soaking, autoclaving, dehulling and micronization followed by air classification have been proposed to improve the nutritional value of peas for poultry (Igbasan and Guenter, 1996; Laudadio *et al.*, 2012; Frikha *et al.*, 2013). These various processing techniques tended to reduce the content of ANF in peas but not efficient enough to be practically used in poultry feed production.

Fermentation processes have been used for centuries in human food production. Fermentation of legumes could eliminate ANF, for example tannins, TI, α -galactosides and modify AA profile by microbial synthesis and breakdown (Ouoba *et al.*, 2003; Ouoba *et al.*, 2007; Gefrom *et al.*, 2013). Moreover, fermentation of an ingredient could provide a high number of beneficial microorganisms with probiotic effects on the microbiology and morphology of the gastrointestinal tract (GIT). Fermentation of rapeseed meal (RSM) could reduce glucosinolate (Chiang *et al.*, 2010) and phytic acid, while increased CP (Nair and Duvnjak, 1990).

One of the major beneficial impacts of fermentation processes on ANF seems to be through enzymatic digestion. Using enzymes as feed additives in poultry diets could inactivate or/and eliminate certain ANF, improve the digestive process and availability of nutrients that were physically or chemically sequestered by ANF substances (Bedford, 2000). The retention time of feed in the GIT of poultry, especially broilers, is very short (~2 to 4 h) and can vary depending on chemical and physical characteristics (i.e. particle size, feed form, NSP concentration, etc.) of the feed. On the other hand, the optimum pH of most exogenous enzymes is between 4 and 6. Consequently, because of the GIT pH and the fact that enzymes can be subjected to hydrolysis by endogenous proteolytic enzymes in the GIT, the degradation activity of exogenous enzymes seems mainly limited to the crop, proventriculus and gizzard (Ravindran, 2013). The short retention time of digesta in the proximal GIT (60 to 90 min) and the wide range of pH which feed encounters along the poultry GIT limits the efficiency of exogenous enzymes (Svihus *et al.*, 2002; Ravindran, 2013).

Fermentation with probiotic microorganisms and treatment of peas for a certain period of time with appropriate exogenous enzymes in a moist environment may be an effective approach to improve the nutritional quality of peas. Despite promising nutritional effects of fermentation and enzymatic treatment of plant materials, there is little information available assessing the effects of these technological processes on nutritional quality of pea as well as on performance and nutrient digestibility in broilers. Therefore, the present study was conducted to

investigate the effects of different inclusion levels of native, fermented or enzymatically treated pea products on performance and nutrient digestibility in broilers.

Material and methods

A commercial batch of Madonna pea (*Pisum sativum L.*) was hammer milled as full seeds (including hulls) to 2 mm screen size. The ground pea (909 g/kg DM) was used for the continuous production of pea dough (500 g/kg DM) during fermentation and enzymatic treatment as well as for the animal trials, as native pea in the diets.

Solid state fermentation

A commercial probiotic, GalliPro[®] (EU authorized probiotic to be used in broiler feed) containing 1.60×10^9 spores/g of *Bacillus subtilis* (Chr. Hansen, Denmark) was employed for the solid state fermentation (SSF) process. In the fermentation process the used water for dough production was inoculated with the probiotic spore, resulting in final concentration of $\sim 2.57 \times 10^8$ *B. subtilis* spores/kg pea. The dough was pumped in internal cathodic protected (ICP) tanks and incubated in temperature controlled containers (30°C) for 48 h.

Treatment with exogenous enzymes

For enzymatic treatment, the used water for dough production contained three different commercial exogenous enzymes plus organic acids mixture (10 ml/kg pea dough; 8 ml lactic acid and 2 ml acetic acid). The acidification of water, resulting in a final pH of 4.5 in the dough, was to inhibit the growth of pea associated microorganisms and uncontrolled fermentation of pea by its associated microorganisms (spoilage) during the incubation time. Three commercial enzymes used for enzymatic treatment process were 0.1 g/kg pea AlphaGal[™] (Kerry EMEA, USA) containing an α -galactosidase (enzyme activity: 115 μ /kg pea), 0.2 g/kg pea RONOZYME[®] ProAct (DSM, Switzerland) containing a protease (enzyme activity: 15 000 μ /kg pea), and 0.2 g/kg pea RONOZYME[®] VP (DSM) containing a blend of β -glucanase and pectinases (enzyme activities: 10 μ /kg pea). After filling in ICP tanks, the prepared dough was incubated for 24 h at 30°C. The effects of fermentation and enzymatic treatment on nutrients and ANF composition of pea products are presented in Tables 1 and 2, respectively.

Drying and milling of processed pea

After the processes, the wet materials were simultaneously dried and ground using an Ultra-Rotor dryer mill (Ultra-Rotor Type U III a, Jäckering Mühlen- und Nahrungsmittelwerke GmbH, Hamm, Germany). The dryer, with generating of a very high turbulence and using of heated air (6000 m³/h), simultaneously milled and dried the material. The drying process lasted <3 s (throughput of 160 kg dry product per h). The maximum product temperature was below 75°C. In total, seven samples from dried products were analyzed concerning their particle size via laser particle size analyzer (Type 930; Cilas S.A., Orléans, France). The mean particle size of the final products

Table 1 Analyzed nutrient composition of pea products

	Native pea ¹	Fermented pea ²	Enzymatically treated pea ²
Nutrient composition (g/kg DM)			
Crude fat	12.1	6.5	6.5
CP	228	238	230
Starch	442	431	434
Ala	9.9	9.9	9.6
Arg	19.1	16.4	18.6
Asp	25.2	24.0	24.6
Cys	5.3	5.2	4.9
Glu	23.9	22.7	22.9
Gly	9.7	9.7	9.5
His	8.1	7.9	7.8
Ile	8.6	8.8	8.7
Leu	15.8	15.8	15.5
Lys	16.6	16.1	16.2
Met	2.6	2.6	2.4
Orn	0.0	2.4	0.0
Phe	10.9	10.9	11.1
Pro	9.9	10.2	9.1
Ser	11.9	11.0	11.4
Thr	9.0	8.9	8.7
Tyr	6.9	6.4	7.1
Val	9.9	10.0	9.9
Total AA	206	205	201
Minerals and trace elements			
Ca (g/kg DM)	0.73	0.90	0.82
K (g/kg DM)	7.60	8.18	8.16
Na (g/kg DM)	0.265	0.230	0.317
P (g/kg DM)	3.17	3.29	3.24
Cu (mg/kg DM)	5.5	6.5	6.5
Fe (mg/kg DM)	42	54	58
Mg (g/kg DM)	1.27	1.21	1.21
Mn (mg/kg DM)	10	11	10
Zn (mg/kg DM)	23	26	28
Bacterial metabolites (µmol/g as fed)			
Acetic acid	2.2	17.8	56.1
Propionic acid	ND	ND	ND
<i>i</i> -Butyric acid ³	ND	ND	ND
<i>n</i> -Butyric acid ³	ND	ND	ND
<i>i</i> -Valeric acid ³	0.8	4.8	1.4
<i>n</i> -Valeric acid ³	14.8	8.6	23.9
L-Lactate	0.52	176	184
D-Lactate	0.05	173	2.0
Ammonium	0.05	32	1.7

ND = not detected.

¹Dry matter of native pea: 909 g/kg.²Dry matter of fermented and enzymatically treated pea: 921 g/kg.³Branched chain volatile fatty acids.

was $\sim 55 \pm 5.2 \mu\text{m}$. The DM content of the final products was about 921 g/kg.

Animals and experimental design

The experimental protocol was approved by the State Office of Health and Social Affairs Berlin (LAGeSo Reg. No. 114-G 0203/14).

Nine starter (days 1 to 21) and nine grower (days 22 to 35) diets were formulated by supplying 10%, 20% and 30% of the required CP with three different pea products tested (native, fermented and enzymatically treated peas). The diets

were formulated to be isocaloric and isonitrogenous for each phase and, meet or exceed the recommendations of the Society of Nutritional Physiology (GfE, 1999). The grower diet contained 5 g titanium dioxide per kg feed (Sigma Aldrich, St. Louis, MO, USA) as an indigestible marker to allow for the determination of ileal apparent nutrient digestibility. The compositions of the starter and grower experimental diets are shown in Tables 3 and 4, respectively.

One thousand and eighty, 1-day-old male broiler chicks (Cobb 500), were randomly allocated into 72 pens (2.20 × 1.80 m) with a softwood shaving floor. Using a completely randomized design with a 3 × 3 factorial arrangement of treatments, the nine different diets were randomly assigned to birds within pens (eight replicate-pens per diet).

The diets were offered in mash. The experiment lasted 35 days. The temperature was 33°C for the first 7 days of the experiment, after which the temperature was gradually reduced by 3°C per week until reaching 24°C. The lighting program consisted of full time light for the first 3 days and 20 h of light until day 7 and 16 h of light thereafter. All birds had access to the experimental diets and water *ad libitum*.

Performance measurements

BWs of the chicks were recorded at day 1 and weekly during the experiment. Feed intake (FI) was recorded weekly and feed conversion ratio (FCR) was calculated. The birds were healthy in the entire experiment and the total mortality was about 2%. Performance variables are presented in Table 5.

Nutrient digestibility and relative organ size

At the end of the experiment, nine birds per pen were randomly selected, stunned and killed by exsanguination. The ileum was dissected from Meckel's diverticulum to the ileo-caeco-colic junction and the digesta was collected from the distal 2/3 for the apparent ileal digestibility (AID) determinations. The digesta of all the birds within each pen were pooled and immediately frozen (−80°C) until further analysis. The pooled digesta was freeze dried before chemical analysis.

The following formula was used for AID calculation:

$$\text{AID of nutrient} = 1 - \left[\frac{(\text{concentration of marker in feed} / \text{concentration of marker in ileum}) \times (\text{concentration of nutrient in ileum} / \text{concentration of nutrient in feed}) \right]$$

The AID of nutrients are presented in Table 6.

Another two birds per pen were weighed, and killed by exsanguination. Carcasses were dissected and the proventriculus, gizzard, duodenum, jejunum, ileum, cecum and pancreas were removed and their empty weights were recorded. Organ size was expressed as a percentage of live BW (Table 7).

Table 2 Anti-nutritional factors of pea products (mean \pm SD)

Anti-nutritional factors	Native pea	Fermentation		Enzymatic treatment	
		Fermented pea	Alteration by fermentation ¹	Enzymatically treated pea	Alteration by treatment ¹
Insoluble-NSP (g/100 g DM) ²	11.3 \pm 0.10	11.1 \pm 0.08	-1.90%	9.9 \pm 0.11	-12.60%
Soluble-NSP (g/100 g DM) ²	0.91 \pm 0.02	1.06 \pm 0.00	+15.40%	1.01 \pm 0.01	+11.00%
Resistant starch (g/100 g DM) ²	3.25 \pm 0.02	0.73 \pm 0.00	-77.50%	0.81 \pm 0.00	-75.10%
Total dietary fiber ³ (g/100 g DM) ²	18.1 \pm 0.11	16.2 \pm 0.05	-10.80%	14.8 \pm 0.12	-18.00%
Trypsin inhibitor activity (TIU) ⁴ (mg/g DM) ²	0.67 \pm 0.01	0.23 \pm 0.01	-65.70%	0.50 \pm 0.01	-25.40%
Raffinose equivalents (mol/g DM) ⁵	97.9 \pm 4.44	30.7 \pm 4.97	-68.60%	48.9 \pm 10.50	-50.00%
Phytic acid (g/100 g DM) ⁵	0.92 \pm 0.02	0.77 \pm 0.01	-16.40%	0.47 \pm 0.01	-49.20%

NSP = non-starch polysaccharides.

¹Compared with native pea.

²Data were obtained from duplicate measurements and are stated as mean value \pm standard deviation.

³Cellulose, hemicellulose, lignin, gums, modified celluloses, mucilages, ligosaccharides, and pectins and associated minor substances, such as waxes, cutin and suberin.

⁴Trypsin inhibitor activity unit.

⁵Data were obtained from triplicate measurements and are stated as mean value \pm standard deviation. Raffinose equivalents represent the number of α -1-6-glycosidic bonds that is cleavable in 1 mol of raffinose.

Table 3 Ingredients (g/kg unless noted) and analyzed nutrient composition of the starter (1 to 21 days) diets¹

Pea	Native			Fermented			Enzymatically treated		
	10%	20%	30%	10%	20%	30%	10%	20%	30%
Level of protein requirement supplementation by pea products									
Ingredient (g/kg)									
Pea product	106.0	212.0	318.0	100.6	201.2	301.8	104.4	208.7	313.1
Maize	234.3	154.5	75.5	243.2	170.3	98.2	237.4	159.2	83.1
Wheat	200.0	200.0	200.0	200.0	200.0	200.0	200.0	200.0	200.0
Soybean meal (CP 440 g/kg)	329.0	288.7	249.5	325.8	285.5	244.5	327.0	288.0	247.8
Soybean oil	82.0	91.5	99.6	81.7	90.0	98.0	82.4	91.0	98.8
Premix (containing 330 g/kg NaCl) ²	12.0	12.0	12.0	12.0	12.0	12.0	12.0	12.0	12.0
Monocalcium phosphate	14.8	15.8	16.2	14.8	15.5	16.2	15.0	15.8	16.2
Limestone	14.4	14.3	14.2	14.4	14.2	14.2	14.3	14.2	14.0
L-Lysine-HCL	2.6	4.2	5.9	2.6	4.3	6.0	2.6	4.2	5.9
D,L-Methionine	3.4	4.3	5.1	3.4	4.3	5.1	3.4	4.2	5.1
L-Threonine	1.2	2.2	3.2	1.2	2.2	3.2	1.2	2.2	3.2
L-Tryptophan	0.3	0.5	0.8	0.3	0.5	0.8	0.3	0.5	0.8
TiO ₂ ³	-	-	-	-	-	-	-	-	-
Analyzed nutrient composition (g/kg)									
CP	232	221	229	225	224	223	223	221	222
Crude fat	101	113	120	103	107	111	104	109	117
Starch	265	241	259	253	267	248	259	243	267
P	7.3	7.5	7.3	7	7.1	7.6	7.2	7.3	7.2
Ca	9.2	9.1	9.3	9.5	8.8	9.4	9.3	9	9.2
Na	1.9	1.9	2.0	2.0	1.9	1.8	2.0	1.9	2.0
Ash	59	59	58	59	59	58	59	59	58
Calculated									
AME _N (MJ/kg) ⁴	12.57	12.57	12.57	12.57	12.57	12.57	12.57	12.57	12.57

¹As-fed basis.

²Contents per kg diet: 4800 IU vitamin A; 480 IU vitamin D₃; 96 mg vitamin E (α -tocopherole acetate); 3.6 mg vitamin K₃; 3 mg vitamin B₁; 3 mg vitamin B₂; 30 mg nicotinic acid; 4.8 mg vitamin B₆; 24 μ g vitamin B₁₂; 300 μ g biotin; 12 mg calcium pantothenic acid; 1.2 mg folic acid; 960 mg choline chloride; 60 mg Zn (zinc oxide); 24 mg Fe (iron carbonate); 72 mg Mn (manganese oxide); 14.4 mg Cu (copper sulfate-pentahydrate); 0.54 mg I (calcium iodate); 0.36 mg Co (cobalt- (II)-sulfate-heptahydrate); 0.42 mg Se (sodium selenite); 1.56 g Na (sodium chloride); 0.66 g Mg (magnesium oxide).

³Indigestible marker (Sigma Aldrich, St. Louis, MO, USA).

⁴Nitrogen-corrected apparent metabolizable energy estimated from chemical composition (based on the EU Regulation – Directive 86/174/EEC): 0.1551 \times % CP + 0.3431 \times % crude fat + 0.1669 \times % starch + 0.1301 \times % total sugar.

Chemical analysis

Basic composition analysis were conducted, using standard procedures (Naumann and Bassler, 2004). A commercial enzymatic test (Starch UV-Test; R-Biopharm, Darmstadt, Germany)

was applied to determine starch content (Naumann and Bassler, 2004). The P content was measured using the ammonium vanadate/molybdate method (Gericke and Kurmies, 1952). Other minerals were analyzed by using an atomic absorption

Table 4 *Ingredients (g/kg unless noted) and analyzed nutrient composition of the grower (22 to 35 days) diets¹*

Pea	Native			Fermented			Enzymatically treated		
	10%	20%	30%	10%	20%	30%	10%	20%	30%
Level of protein requirement supplementation by pea products									
Ingredient (g/kg)									
Pea product	89.1	178.1	267.2	84.5	169.1	253.6	87.7	175.3	263.0
Maize	280.7	210.0	146.7	287.2	227.1	166.8	282.6	218.3	153.4
Wheat	300.0	300.0	300.0	300.0	300.0	300.0	300.0	300.0	300.0
Soybean meal (CP 440 g/kg)	215.7	186.3	150.4	214.5	180.2	146.0	215.4	182.3	148.8
Soybean oil	64.2	72.0	78.5	63.6	70.0	76.6	64.0	70.6	77.5
Premix (containing 330 g/kg NaCl) ²	12.0	12.0	12.0	12.0	12.0	12.0	12.0	12.0	12.0
Monocalcium phosphate	11.9	12.1	12.7	11.8	12.1	12.7	11.9	12.1	12.8
Limestone	11.0	11.1	11.0	11.0	11.1	10.8	11.0	11.1	10.9
L-Lysine-HCL	4.3	5.6	7.0	4.3	5.7	7.1	4.3	5.6	7.1
D,L-Methionine	3.2	3.9	4.6	3.2	3.9	4.5	3.2	3.9	4.6
L-Threonine	2.1	2.9	3.7	2.1	2.8	3.7	2.1	2.8	3.7
L-Tryptophan	0.8	1.0	1.2	0.8	1.0	1.2	0.8	1.0	1.2
TiO ₂ ³	5.0	5.0	5.0	5.0	5.0	5.0	5.0	5.0	5.0
Analyzed nutrient composition (g/kg)									
CP	188	189	185	186	188	187	187	187	190
Crude fat	90	88	96	86	88	88	86	88	92
Starch	341	339	340	325	313	346	369	345	323
P	6.1	5.8	5.7	6.0	6.1	6.1	6.0	5.9	6.0
Ca	7.7	7.2	7.1	7.5	7.4	7.4	7.7	7.7	7.7
Na	1.8	1.6	1.9	1.9	1.8	1.8	1.9	1.9	2.0
Ash	54	52	52	54	53	52	54	54	53
Calculated AME _N (MJ/kg) ⁴	12.65	12.65	12.65	12.65	12.65	12.65	12.65	12.65	12.65

¹As-fed basis.

²Contents per kg diet: 4800 IU vitamin A; 480 IU vitamin D₃; 96 mg vitamin E (α-tocopherole acetate); 3.6 mg vitamin K₃; 3 mg vitamin B₁; 3 mg vitamin B₂; 30 mg nicotinic acid; 4.8 mg vitamin B₆; 24 μg vitamin B₁₂; 300 μg biotin; 12 mg calcium pantothenic acid; 1.2 mg folic acid; 960 mg choline chloride; 60 mg Zn (zinc oxide); 24 mg Fe (iron carbonate); 72 mg Mn (manganese oxide); 14.4 mg Cu (copper sulfate-pentahydrate); 0.54 mg I (calcium iodate); 0.36 mg Co (cobalt-(II)-sulfate-heptahydrate); 0.42 mg Se (sodium selenite); 1.56 g Na (sodium chloride); 0.66 g Mg (magnesium oxide).

³Indigestible marker (Sigma Aldrich, St. Louis, MO, USA).

⁴Nitrogen-corrected apparent metabolizable energy estimated from chemical composition (based on the EU Regulation – Directive 86/174/EEC): 0.1551 × % CP + 0.3431 × % crude fat + 0.1669 × % starch + 0.1301 × % total sugar.

Table 5 *Effect of the experimental diets on the performance variables (mean¹ values)*

	Type of the pea products			Level ²			SEM	P value		
	Native	Fermented	Enzymatically treated	10%	20%	30%		Type ³	Level	Type × level
BWG (1 to 21 days)	690	685	696	718 ^a	691 ^{ab}	662 ^b	8.3	Ns ⁴	0.025	Ns
FI (1 to 21 days)	900	881	879	939 ^a	883 ^{ab}	838 ^b	11.7	Ns	0.002	Ns
FCR (1 to 21 days)	1.31	1.29	1.26	1.31	1.28	1.27	0.009	Ns	Ns	Ns
BWG (22 to 35 days)	1119	1078	1093	1133 ^a	1117 ^a	1041 ^b	8.9	Ns	<0.001	Ns
FI (22 to 35 days)	1783 ^a	1712 ^b	1699 ^b	1795 ^a	1753 ^a	1645 ^b	14.7	0.010	<0.001	Ns
FCR (22 to 35 days)	1.59 ^a	1.59 ^a	1.55 ^b	1.59	1.57	1.58	0.006	0.017	Ns	Ns
BWG (1 to 35 days)	1809	1763	1789	1851 ^a	1808 ^a	1702 ^b	14.1	Ns	<0.001	Ns
FI (1 to 35 days)	2683 ^a	2593 ^{ab}	2577 ^b	2734 ^a	2636 ^{ab}	2483 ^b	21.9	0.039	<0.001	Ns
FCR (1 to 35 days)	1.48 ^a	1.47 ^a	1.44 ^b	1.48	1.46	1.46	0.005	0.002	Ns	Ns

SEM = pooled standard error of mean; FCR = feed conversion ratio (g of feed intake/g of BW gain); FI = feed intake (g); BWG = BW gain (g)

^{a,b}Means of each main factors with different superscripts in a row differ significantly ($P \leq 0.05$).

¹Data are means of eight replicate pens with 15 birds per pen.

²Level of protein requirement supplementation by pea products.

³Type of the pea products.

⁴Not significant ($P > 0.05$). Differences were considered significant at $P \leq 0.05$.

spectrophotometer (AAS vario®; Analytik Jena, Jena, Germany). TiO₂ content was measured using the method described by Short *et al.* (1996). The AA analyses were conducted using a Biochrom 20 Plus amino acid analyzer (Amersham Pharmacia

Biotech, Piscataway, NJ, USA) using standard method (VDLUF, 2003).

The soluble and insoluble NSP were determined using Englyst and Cummings (1984) method. The American

Table 6 Effect of the experimental diets on the apparent ileal digestibility (AID) of nutrient (mean¹ values) in broilers (day 35)

	Type of the pea products			Level ²			SEM	P value		
	Native	Fermented	Enzymatically treated	10%	20%	30%		Type ³	Level	Type × level
Ala	0.82	0.82	0.81	0.81	0.82	0.82	0.005	Ns ⁴	Ns	Ns
Arg	0.89	0.89	0.89	0.88	0.89	0.89	0.003	Ns	Ns	Ns
Asp	0.80	0.80	0.80	0.79	0.80	0.81	0.004	Ns	Ns	Ns
Cys	0.75	0.75	0.74	0.74	0.75	0.75	0.006	Ns	Ns	Ns
Glu	0.86	0.86	0.85	0.85	0.86	0.86	0.003	Ns	Ns	Ns
Gly	0.78	0.79	0.78	0.78	0.78	0.79	0.005	Ns	Ns	Ns
His	0.84	0.84	0.83	0.83	0.84	0.84	0.004	Ns	Ns	Ns
Ile	0.83	0.83	0.83	0.83	0.82	0.83	0.006	Ns	Ns	Ns
Leu	0.83	0.84	0.83	0.83	0.83	0.84	0.005	Ns	Ns	Ns
Lys	0.89	0.90	0.89	0.88 ^b	0.89 ^{ab}	0.91 ^a	0.003	Ns	0.009	Ns
Met	0.95	0.95	0.95	0.94 ^b	0.95 ^{ab}	0.96 ^a	0.002	Ns	0.007	Ns
Phe	0.85	0.86	0.85	0.85	0.85	0.86	0.005	Ns	Ns	Ns
Pro	0.82	0.83	0.83	0.82	0.82	0.83	0.005	Ns	Ns	Ns
Ser	0.80	0.80	0.80	0.80	0.80	0.81	0.004	Ns	Ns	Ns
Thr	0.82	0.82	0.82	0.80 ^b	0.82 ^{ab}	0.84 ^a	0.005	Ns	0.011	Ns
Tyr	0.83	0.84	0.84	0.83	0.83	0.84	0.005	Ns	Ns	Ns
Val	0.80	0.80	0.80	0.80	0.80	0.81	0.007	Ns	Ns	Ns
Total AA	0.83	0.84	0.83	0.83	0.83	0.84	0.004	Ns	Ns	Ns
CP	0.82	0.82	0.81	0.81	0.82	0.83	0.004	Ns	Ns	Ns
CF	0.93	0.93	0.93	0.93	0.93	0.93	0.004	Ns	Ns	Ns
Starch	0.92	0.96	0.96	0.96	0.94	0.93	0.004	<0.001	<0.001	0.018
P	0.60	0.62	0.60	0.61	0.62	0.61	0.010	Ns	Ns	Ns
Ca	0.38	0.37	0.38	0.38	0.37	0.39	0.010	Ns	Ns	Ns
K	0.85	0.85	0.83	0.84	0.85	0.84	0.004	Ns	Ns	Ns

SEM = pooled standard error of mean; AA = amino acids; CF = crude fat.

^{a,b}Means of each main factors with different superscripts in a row differ significantly ($P \leq 0.05$).

¹Data are means of eight replicate pens. Pooled digesta of nine birds per pen.

²Level of protein requirement supplementation by pea products.

³Type of the pea products.

⁴Not significant ($P > 0.05$). Differences were considered significant at $P \leq 0.05$.

Table 7 Effect of the experimental diets on relative organ weight (day 35) of broilers¹ (mean² values)

	Type of the pea products			Level ³			SEM	P value		
	Native	Fermented	Enzymatically treated	10%	20%	30%		Type ⁴	Level	Type × level
Pancreas	0.18	0.19	0.18	0.19	0.18	0.18	0.004	Ns ⁵	Ns	Ns
Proventriculus	0.29	0.30	0.27	0.29	0.28	0.30	0.006	Ns	Ns	Ns
Gizzard	1.66 ^a	1.41 ^b	1.40 ^b	1.46	1.48	1.53	0.031	<0.001	Ns	Ns
Duodenum	0.71 ^a	0.67 ^{ab}	0.64 ^b	0.67	0.67	0.68	0.011	0.016	Ns	Ns
Jejunum	1.18 ^a	1.11 ^{ab}	1.08 ^b	1.12	1.15	1.10	0.018	0.050	Ns	Ns
Ileum	0.88 ^a	0.80 ^b	0.79 ^b	0.84	0.80	0.83	0.014	0.014	Ns	Ns
Small intestine	2.78 ^a	2.59 ^{ab}	2.50 ^b	2.63	2.63	2.62	0.036	0.004	Ns	Ns
Cecum	0.30	0.29	0.30	0.29	0.29	0.31	0.008	Ns	Ns	Ns

SEM = pooled standard error of mean.

^{a,b}Means with different superscripts in a row differ significantly ($P \leq 0.05$).

¹[Empty organ weight (g)/live BW (g)] × 100.

²Data are means of eight replicate pens (two birds per pen).

³Level of protein requirement supplementation by pea products.

⁴Type of the pea products.

⁵Not significant ($P > 0.05$). Differences were considered significant at $P \leq 0.05$.

Association of Cereal Chemists (2003) methods were used to determine RS (AACC 32-40.01) and total dietary fiber (TDF; AACC method 32-25 – Uppsala method). Trypsin inhibitor

activity (TIA) was measured according to Kakade *et al.* (1974) method. The α -galactosides content (in pea mainly raffinose, stachyose and verbascose) was determined via enzymatic

assay (raffinose/ β -galactose Assay Kit; Megazyme International, Bray, Ireland), presented as raffinose equivalent (RE). Phytate was analyzed using phytate (total P) assay kit (Megazyme International).

Determination of bacterial metabolites

Determination of short chain fatty acids was performed by gas chromatography (Agilent Technologies 6890N coupled with an auto sampler G2614A and an auto injector G2613A; Santa Clara, CA, USA) using standard procedure (Schäfer, 1995). The column was an Agilent 19095N-123 HP-INNOWAX polyethylene glycol.

Following standard procedure, D- and L-lactate were determined by HPLC, using an Agilent 1100 system with a Phenomenex C18 (4.0 \times 2.0 mm) guard column followed by a Phenomenex Chirex 3126 (D)-penicillamine column (150 \times 4.6 mm) (Agilent Technologies). The UV detector wavelength was 253 nm and the column temperature was 35°C. The carrier was CuSO₄ in a gradient from 0.5 to 2.5 mmol/l with a flow rate of 1 ml/min at 35°C and the injection volume was 20 μ l.

Ammonia was quantified using the Berthelot reaction assay. Briefly, 20 μ l of the sample was chlorinated with 100 μ l of 0.2% alkaline hypochloride (Sigma Aldrich, Deisenhofen, Germany) to convert NH₃ to chloramine (NH₂Cl) following reaction with thymol to N-chloro-2-isopropyl-5-methylchinonmonoimin and further to indophenol using 100 μ l of 5% phenol nitroprusside (Sigma Aldrich). After incubation for 100 min in microtitration plates at room temperature, a photometric measurement was carried out at 620 nm with a Tecan microtiterplate reader (Tecan Austria GmbH, Grödig, Austria).

Statistical analysis

Data were subjected to ANOVA using the GLM procedure of SPSS 19.0 (SPSS Inc., Chicago, IL, USA) as a 3 \times 3 factorial arrangement of treatments that included three pea products (native, fermented or enzymatically treated pea) and three different inclusion levels (10%, 20% or 30% of diet's CP) as the main effects and their interactions. Treatment means were separated by the Tukey least significant difference *post hoc* test at $P \leq 0.05$ statistical level. Replicate-pen was the experimental unit for all variables measured.

Results

Influence of technological processing on nutrients and anti-nutritional factors of the pea products

Except a reduction in crude fat (CF) and a negligible increase in CP, no other notable changes for the main nutrients like AA and starch were observed for the both processed pea products. Both processes caused a minor increase in the concentration of P, Ca, K, Fe and partially Zn. The Na content of enzymatically treated pea was higher than native and fermented ones.

In accordance with pH value of fermented pea, which decreased from 6.34 down to 4.37 within 48 h incubation

time, the concentration of acetic acid, L-lactate, D-lactate and ammonium in fermented pea were considerably higher than native pea. As expected, the pH value of the enzymatically treated pea did not vary with time (4.54 ± 0.14).

The RE content of pea was reduced by 69% in the fermented end product (30.7 mol/g) and by 50% in the enzymatically treated one (48.9 mol/g). Fermentation decreased TIA in the pea (0.67 TIU mg/g) to approximately one third (0.23 TIU mg/g), whereas TIA in the enzymatically treated pea was ~25% (0.50 TIU mg/g) lower than the native pea. Enzymatic treatment caused a 49% reduction in phytic acid concentration, whereas the reduction by fermentation process was only of 16% (0.92 g phytate/100 g native pea).

Both processes caused a slight increase in S-NSP and a slight reduction in I-NSP and TDF, with reductions being more pronounced for the enzymatically treated pea. Moreover, a remarkable decrease in RS was observed in both processed peas (more than 75% reduction).

Broiler performance

Increasing the level of dietary peas in the diets up to 30% reduced BWG and FI of broilers ($P \leq 0.05$). Pea processing decreased FI after day 21 ($P \leq 0.05$), whereas broilers fed enzymatically treated pea had the best FCR for the growing and entire experimental period ($P \leq 0.05$). At the end of the experiment, FI of birds fed enzymatically treated pea diets was considerably lower than those received native pea diets ($P \leq 0.05$).

Apparent ileal nutrient digestibility and relative organ size

The AID of Thr, Lys and Met at 30% inclusion level was higher than 10% inclusion group ($P \leq 0.05$). The interaction between type of the pea products and inclusion levels of peas was only significant for the AID of starch ($P \leq 0.05$). Chicken fed 30% native pea diet had the lowest AID of starch (0.89) among all the nine groups and chicken fed 10% fermented and enzymatically treated pea diets (0.97 and 0.97, respectively) had higher AID of starch compared with 20% and 30% native pea groups (0.93 and 0.89, respectively). In native pea groups, AID of starch for 20% (0.93) and 30% (0.89) native pea diets were similar but 10% native pea diet (0.95) showed similar AID to 20% and higher AID than 30% one ($P \leq 0.05$). In fermented pea groups (0.97, 0.95 and 0.95, respectively) as well as enzymatically treated groups (0.97, 0.96 and 0.96, respectively), all three levels had identical AID of starch.

The size of gizzard, duodenum, jejunum and ileum, were significantly lower for birds fed enzymatically treated pea compared with those received native pea ($P \leq 0.05$). The size of gizzard and ileum in broilers received fermented pea diets followed the course observed for those fed enzymatically treated peas ($P \leq 0.05$).

Discussion

Fermentation processes to improve nutritional quality of legumes, particularly soybean, have been studied for long (Feng *et al.*, 2007). The application of exogenous enzymes in

poultry diets to overcome the negative impacts of ANF substances and improve the nutritional quality of feed is a common practice in poultry feed production. The main limits for optimal enzyme responses in poultry nutrition seem to be the retention time and variable pH in the different segments of the GIT (Adeola and Cowieson, 2011). Selective treatment by enzymes and also fermentation of SBM could eliminate the allergenic proteins and soluble carbohydrates, and led to a reduction in fiber, TI, oligosaccharides and lectins (Berrocoso *et al.*, 2013), consequently, they could improve growth performance and nutrient digestibility in poultry (Frikha *et al.*, 2013). Subjecting pea to fermentation with probiotic microorganisms or treatment with appropriate exogenous enzymes, may be a sound alternative to improve feeding value of plant protein sources to replace SBM in poultry diet.

Fermentation of African locust bean (*Parkia biglobosa*) at pH 7.5 to 9 by different *Bacillus subtilis* subspecies increased essential AA and unsaturated fatty acids, for example linoleic and linolenic acids content, and also decreased oligosaccharides (Ouoba *et al.*, 2003 and 2007). Fermentation of RSM with *Lactobacillus fermentum* and *B. subtilis* increased CP, Lys, sulfur AA and considerably decreased isothiocyanates (from 108.7 to 13.1 mmol/kg) (Xu *et al.*, 2012). In the present study none of the two processes modified the AA content of the peas, except a reduction in CF and a slight increase in CP. Since the alteration in fat content was also presented in the not fermented samples (enzymatically treated sample), it is very likely that fat reduction was due to the lipase activity of peas. This is in accordance with other study which reported a distinct reduction of lipids and an increase in lipase activity during fermentation of soybean by its associated (native flora – no bacteria added) microorganisms (Ruiz-Teran and Owens, 1996). In the present study, the reason for observed minor increases in the concentration of analyzed minerals by the processes is not clear. Fermentation and enzymatic treatment of pea in lab scale did not cause any change in ash content of the final products (unpublished data). Thus, considering the amount of water used for pea dough production (50%), this minor increases might be explained by mineral content of the water in the production site (Jäckering Mühlen- und Nahrungsmittelwerke GmbH, Hamm, Germany) and also wear of metallic pieces in the Ultra-Rotor dryer. Increase in mineral content of poultry feed due to the wear of metallic pieces in hydrothermal processing machines has been reported before (Goodarzi Borojjeni *et al.*, 2016). Both types of pea processing resulted in a remarkable decrease in RS. The slight reductions in I-NSP and TDF of the processed peas were more pronounced for the enzymatically treated pea than the fermented one. Enzymatic treatment reduced TIA and phytic acid by 25% and 49%, while these were reduced by 66% and 16% in the fermented pea. The RE was reduced by 50% and 69% in the enzymatically treated and fermented peas, respectively. The results are in accordance with recent study which showed that SSF of SBM with *B. subtilis* decreased the TIA of SBM up to 95% (Teng *et al.*, 2012). The changes in phytic acid, RE, I-NSP and TDF might be explained by activation of exogenous/microbial enzymes and pea's endogenous enzymes.

Furthermore, the acidic pH value in the processed peas might have induced an increase in activity of endogenous amylase and phytase (Selle *et al.*, 2000; Adeola and Cowieson, 2011). The drastic pH drop during the SSF process seemed to be due to the accumulation of lactic and acetic acids, caused by the metabolic activity of native and added microorganisms (Ying *et al.*, 2009). The high concentration of acetic and L-lactic acids in the enzymatically treated pea was mainly due to the addition of organic acids during the processing. The increase of ammonium, valeric and D-lactic acids by enzymatic treatment could be because of activity and proliferation of the pea's native microflora as well as existent microorganisms in the production environment, which were able to survive and maintain their metabolic activity at the created acidic pH (4.5).

The information available regarding the effects of enzymatic treatment on nutrient and ANF composition of peas is scarce. In a broiler study, enzymatic treatment of pea with several carbohydrase enzymes (i.e. cellulase, pectinase, xylanase, glucanase, galactanase, and mannanase) and their combinations reduced arabinose, xylose, galactose, glucose, uronic acids and total NSP. These reductions were more pronounced when a combination of enzymes was used (Meng *et al.*, 2005). In another broiler study, soaking of barley for 24 h at room temperature with a commercial complex (with protease, xylanase and β -glucanase activity) reduced acid extract viscosity as well as soluble, insoluble and total β -glucan concentration in barley (Svihus *et al.*, 1997). Treatment of brewers' spent grain with xylanase for 3 h reduced the concentration of xylose and arabinose by 15% to 30% (Denstadli *et al.*, 2010).

Inclusion of processed peas in the diets reduced FI during the growing period, with birds fed enzymatically treated pea showing the best FCR for the growing (days 22 to 35) and entire experimental period. Treatment of whole barley with a mixture of protease, xylanase and β -glucanase improved BWG and FCR in broiler chicks (Svihus *et al.*, 1997), whereas in another study inclusion of enzymatically treated (with xylanase) brewers' spent grain in broiler diets increased FI with no beneficial impact on BWG and FCR (Denstadli *et al.*, 2010). The observed FCR improvement in the enzymatically treated group might be due to the observed lower FI or/and the lower concentrations of ANF in the enzymatically treated peas as well as the degradation effect of supplemented enzymes on complex nutrients. In an experiment, Chen *et al.* (2009) investigated whether the beneficial effect of fermented feed on broiler growth performance was because of the probiotics *per se* or the fermentation process. Inclusion of probiotic, with similar microflora population to the fermented feed, did not enhance growth performance as much as fermented feed did. The authors explained this observation by degradation effect of fermentation process on complex material and production of beneficial substances for broiler growth and health. In the present study, inclusion of fermented pea in broiler diets did not improve growth performance and nutrient digestibility. The total tract apparent digestibility of DM, energy and Ca were higher in broilers received SSF RSM than in those fed native RSM (Chiang *et al.*, 2010). Chickens fed fermented SBM with

Aspergillus oryzae showed better FI, BWG and FCR compared with those received native SBM (Feng *et al.*, 2007). Broilers fed diets containing 10% fermented RSM (with *L. fermentum*, *Enterococcus faecium*, *Saccharomyces cerevisiae* and *B. subtilis*) had better BWG and FCR compared with those fed diets containing 10% native RSM (Chiang *et al.*, 2010). The BWG and FCR of broilers fed diets containing 15% fermented RSM were worse than those received feed containing 0, 5 and 10% fermented RSM, while the other three groups (0%, 5% and 10%) were similar (Xu *et al.*, 2012).

The interaction between type of the pea products and inclusion levels of peas was significant for AID of starch. Increasing in inclusion level of native pea in broiler diets from 10% to 30% reduced AID of starch. In agreement with the present results, Brenes *et al.* (1993) and Igbasan and Guenter (1996) showed that inclusion of native pea in broiler diets resulted in significant reduction of ileal starch digestibility. This reduction in AID of starch by increasing inclusion level was not observable anymore when fermented and enzymatically treated peas were used in broiler diets. This might be explained by the reduction of RS as well as starch swelling and gelatinization that might have happened during processing and drying. In a review manuscript, it was demonstrated that the temperature threshold for the stabilization of starch gelatinization has a negative correlation with the starch hydration (Abdollahi *et al.*, 2013). For instance, at excess water content (above 40%), the starch gelatinization was stabilized at a temperature between 50°C and 70°C. This temperature increased to above 100°C with <35% water content (Svihus *et al.*, 2005). Processed peas in the present study were incubated in a moist condition (>40% water content) and were dried using a relatively mild thermal and shear treatments (<75°C for <3 s). The applied production and drying conditions might lead to starch swelling and partial gelatinization which may have caused alteration in starch digestibility.

In the present study, the inclusion level of pea products had no remarkable effect on digestibility of AA. The observed increase in the AID of Thr, Lys and Met in the 30% inclusion group seemed to be only due to the higher levels of crystalline Thr, Lys and Met in 30% inclusion diets. Furthermore, 30% inclusion group had lower FI and BWG compared with 10% inclusion group. It has been recommended that peas can be used up to 200 g/kg in broiler diet as a protein source with no negative impact on growth performance (Nalle *et al.*, 2011). In a review paper, the maximum inclusion level of 200 g/kg peas in broiler diets has been recommended (Castell *et al.*, 1996), while in another study the maximum inclusion level of 300 g/kg in broiler diets was suggested (Farrell *et al.*, 1999). Increase in inclusion levels of whole canola/pea mixture (0, 100, 200 or 300 g/kg) in broiler diets linearly declined BWG and curvilinearly FI, while the reduction in FI was most apparent at higher concentrations (Fasina and Campbell, 1997).

It is a well-documented phenomenon that the GIT of broilers adapts quickly to the alterations in diet structure and composition (Svihus, 2011). The lower relative organ weight of

the different gut sections in both processed pea groups, particularly for birds fed enzymatically treated pea, could be explained by observed reduction in ANF, degradation of complex material and better availability of nutrients in the processed peas. It has been reported that a decrease in ANF content of feed could lead to reduction in digesta viscosity, alteration in the gut microbiota and changes in the morphology of the digestive tract (Svihus *et al.*, 1997; Bedford and Cowieson, 2012). In agreement with the present data, broilers fed enzymatically treated barley diets had lower relative gizzard and small intestine weights as well as similar relative cecum and pancreas weights compared with broilers fed native barley diets (Svihus *et al.*, 1997). Gizzard development is directly related to the feed particle size. Fine feed particles in poultry diets coincide with gizzard under-development (Svihus, 2011). In the present study, the Ultra-Rotor dryer mill which has been used to dry fermented and enzymatically treated peas, simultaneously milled and dried the material. Therefore the final particle sizes of the processed peas were considerably smaller than the initial particle size of the ground native pea which has been used for production of the processed peas as well as the native pea diets in the animal trial. As the particle sizes of wheat, corn and other feed ingredients used in broiler diets were similar, it can be speculated that finer particle sizes of fermented and enzymatically treated peas led to poor gizzard development.

Fermentation and enzymatic treatment could improve the nutritional quality of pea by reduction in ANFs. Furthermore, taken into account insignificant interactions between type of the pea products and inclusion levels for variables measured (except for ileal starch digestibility), it can be concluded that inclusion of enzymatically treated pea in broiler diets could improve broiler performance compared with other pea products while, it showed neither positive nor negative impact on nutrients digestibility. However, the mode of action for feed efficiency improvement by inclusion of enzymatically treated pea is not clear enough and needs to be studied further. The present findings indicate the feasibility of these processes, particularly enzymatic treatment, for improving the nutritional quality of pea as a protein source for broiler nutrition.

Acknowledgment

This work has been funded by CORNET through AiF Projekt GmbH (106 EN/1 (OE142/13a)). The authors express deep appreciation to Dr W. Vahjen, Dr T. Roick, Dr J. Jacobi, Dr S. Rohn, A. Kriesten, K. Topp, L. Rod, I. Bebert, C. Schmidt, P. Feige and I. Bayram for technical support during the animal experiment and laboratory analysis.

References

- Abdollahi MR, Ravindran V and Svihus B 2013. Pelleting of broiler diets: an overview with emphasis on pellet quality and nutritional value. *Animal Feed Science and Technology* 179, 1–23.
- Adeola O and Cowieson A 2011. Board-invited review: opportunities and challenges in using exogenous enzymes to improve nonruminant animal production. *Journal of Animal Science* 89, 3189–3218.

- American Association of Cereal Chemists 2003. Methods: 32–25 (dietary fibre), 32-40.01 (resistant starch). American Association of Cereal Chemists, St. Paul, Minnesota, USA.
- Bedford M and Cowieson A 2012. Exogenous enzymes and their effects on intestinal microbiology. *Animal Feed Science and Technology* 173, 76–85.
- Bedford MR 2000. Exogenous enzymes in monogastric nutrition – their current value and future benefits. *Animal Feed Science and Technology* 86, 1–13.
- Berrococo J, Serrano M, Cámara L, López A and Mateos G 2013. Influence of source and micronization of soybean meal on nutrient digestibility and growth performance of weanling pigs. *Journal of Animal Science* 91, 309–317.
- Brenes A, Rotter B, Marquardt R and Guenter W 1993. The nutritional value of raw, autoclaved and dehulled peas (*Pisum sativum* L.) in chicken diets as affected by enzyme supplementation. *Canadian Journal of Animal Science* 73, 605–614.
- Castell A, Guenter W and Igbanan F 1996. Nutritive value of peas for nonruminant diets. *Animal Feed Science and Technology* 60, 209–227.
- Chen K-L, Kho W-L, You S-H, Yeh R-H, Tang S-W and Hsieh C-W 2009. Effects of *Bacillus subtilis* var. *natto* and *Saccharomyces cerevisiae* mixed fermented feed on the enhanced growth performance of broilers. *Poultry Science* 88, 309–315.
- Chiang G, Lu W, Piao X, Hu J, Gong L and Thacker P 2010. Effects of feeding solid-state fermented rapeseed meal on performance, nutrient digestibility, intestinal ecology and intestinal morphology of broiler chickens. *Asian-Australasian Journal of Animal Sciences* 23, 263–271.
- Denstadli V, Westereng B, Biniyam HG, Ballance S, Knutsen SH and Svihus B 2010. Effects of structure and xylanase treatment of brewers' spent grain on performance and nutrient availability in broiler chickens. *British Poultry Science* 51, 419–426.
- Englyst HN and Cummings JH 1984. Simplified method for the measurement of total non-starch polysaccharides by gas-liquid chromatography of constituent sugars as alditol acetates. *Analyst* 109, 937–942.
- Farrell DJ, Perez-Maldonado RA and Mannion PF 1999. Optimum inclusion of field peas, faba beans, chick peas and sweet lupins in poultry diets. II. Broiler experiments. *British Poultry Science* 40, 674–680.
- Fasina Y and Campbell G 1997. Whole canola/pea and whole canola/canola meal blends in diets for broiler chickens. 2. Determination of optimal inclusion levels. *Canadian Journal of Animal Science* 77, 191–195.
- Feng J, Liu X, Xu Z, Wang Y and Liu J 2007. Effects of fermented soybean meal on digestive enzyme activities and intestinal morphology in broilers. *Poultry Science* 86, 1149–1154.
- Frikha M, Valencia D, de Coca-Sinova A, Lázaro R and Mateos G 2013. Ileal digestibility of amino acids of unheated and autoclaved pea protein concentrate in broilers. *Poultry Science* 92, 1848–1857.
- Gefrom A, Ott E, Hoedtke S and Zeyner A 2013. Effect of ensiling moist field bean (*Vicia faba*), pea (*Pisum sativum*) and lupine (*Lupinus spp.*) grains on the contents of alkaloids, oligosaccharides and tannins. *Journal of Animal Physiology and Animal Nutrition* 97, 1152–1160.
- Gericke S and Kurmies B 1952. Die kolorimetrische Phosphorsäurebestimmung mit Ammonium-Vanadat-Molybdat und ihre Anwendung in der Pflanzenanalyse. *Z Pflanzenernaehr Dueng Bodenkd* 59, 235–247.
- GfE 1999. Empfehlungen zur Energie- und Nährstoffversorgung der Legehennen und Masthühner Broiler. DLG-Verlag, Frankfurt (Main), Germany.
- Goodarzi Borojani F, Svihus B, von Reichenbach HG and Zentek J 2016. The effects of hydrothermal processing on feed hygiene, nutrient availability, intestinal microbiota and morphology in poultry – a review. *Animal Feed Science and Technology* 220, 187–215.
- Igbanan F and Guenter W 1996. The evaluation and enhancement of the nutritive value of yellow-, green- and brown-seeded pea cultivars for unpelleted diets given to broiler chickens. *Animal Feed Science and Technology* 63, 9–24.
- Kakade M, Rackis J, McGhee J and Puski G 1974. Determination of trypsin inhibitor activity of soy products: a collaborative analysis of an improved procedure. *Cereal Chemistry* 51, 376–382.
- Laudadio V, Nahashon S and Tufarelli V 2012. Growth performance and carcass characteristics of guinea fowl broilers fed micronized-dehulled pea (*Pisum sativum* L.) as a substitute for soybean meal. *Poultry Science* 91, 2988–2996.
- Meng X, Slominski B, Nyachoti C, Campbell L and Guenter W 2005. Degradation of cell wall polysaccharides by combinations of carbohydrase enzymes and their effect on nutrient utilization and broiler chicken performance. *Poultry Science* 84, 37–47.
- Nair V and Duvnjak Z 1990. Reduction of phytic acid content in canola meal by *Aspergillus ficuum* in solid state fermentation process. *Applied Microbiology and Biotechnology* 34, 183–188.
- Nalle C, Ravindran V and Ravindran G 2011. Nutritional value of peas (*Pisum sativum* L.) for broilers: apparent metabolisable energy, apparent ileal amino acid digestibility and production performance. *Animal Production Science* 51, 150–155.
- Naumann K and Bassler R 2004. Methodenbuch Band III: Die chemische Untersuchung von Futtermitteln. Neumann-Neudamm, Melsungen, Germany.
- Ouoba L, Cantor M, Diawara B, Traore A and Jakobsen M 2003. Degradation of African locust bean oil by *Bacillus subtilis* and *Bacillus pumilus* isolated from soumbala, a fermented African locust bean condiment. *Journal of Applied Microbiology* 95, 868–873.
- Ouoba LI, Diawara B, Christensen T, Mikkelsen JD and Jakobsen M 2007. Degradation of polysaccharides and non-digestible oligosaccharides by *Bacillus subtilis* and *Bacillus pumilus* isolated from Soumbala, a fermented African locust bean (*Parkia biglobosa*) food Condiment. *European Food Research and Technology* 224, 689–694.
- Ravindran V 2013. Feed enzymes: the science, practice, and metabolic realities. *The Journal of Applied Poultry Research* 22, 628–636.
- Ruiz-Teran F and Owens DJ 1996. Chemical and enzymic changes during the fermentation of bacteria-free soya bean tempe. *Journal of the Science of Food and Agriculture* 71, 523–530.
- Schäfer K 1995. Analysis of short chain fatty acids from different intestinal samples by capillary gas chromatography. *Chromatographia* 40, 550–556.
- Selle P, Ravindran V, Caldwell A and Bryden W 2000. Phytate and phytase: consequences for protein utilisation. *Nutrition Research Reviews* 13, 255–278.
- Short F, Gorton P, Wiseman J and Boorman K 1996. Determination of titanium dioxide added as an inert marker in chicken digestibility studies. *Animal Feed Science and Technology* 59, 215–221.
- Svihus B 2011. The gizzard: function, influence of diet structure and effects on nutrient. *World's Poultry Science Journal* 67, 207–224.
- Svihus B, Newman RK and Newman CW 1997. Effect of soaking, germination, and enzyme treatment of whole barley on nutritional value and digestive tract parameters of broiler chickens. *British Poultry Science* 38, 390–396.
- Svihus B, Hetland H, Choct M and Sundby F 2002. Passage rate through the anterior digestive tract of broiler chickens fed on diets with ground and whole wheat. *British Poultry Science* 43, 662–668.
- Svihus B, Uhlen AK and Harstad OM 2005. Effect of starch granule structure, associated components and processing on nutritive value of cereal starch: a review. *Animal Feed Science and Technology* 122, 303–320.
- Teng D, Gao M, Yang Y, Liu B, Tian Z and Wang J 2012. Bio-modification of soybean meal with *Bacillus subtilis* or *Aspergillus oryzae*. *Biocatalysis and Agricultural Biotechnology* 1, 32–38.
- VDLUF 2003. Handbuch der landwirtschaftlichen Versuchs- und Untersuchungsmethodik, Methodenbuch Band VI-Chemische, physikalische und mikrobiologische Untersuchungsverfahren für Milch Milchprodukte und Molkereihilfsstoffe. VDLUF-Verlag, Darmstadt, Germany.
- Xu F, Zeng X and Ding X 2012. Effects of replacing soybean meal with fermented rapeseed meal on performance, serum biochemical variables and intestinal morphology of broilers. *Asian-Australasian Journal of Animal Sciences* 25, 1734.
- Ying W, Zhu R, Lu W and Gong L 2009. A new strategy to apply *Bacillus subtilis* MA139 for the production of solid-state fermentation feed. *Letters in Applied Microbiology* 49, 229–234.